

# Epistatic interaction between *Arabidopsis FRI* and *FLC* flowering time genes generates a latitudinal cline in a life history trait

Ana L. Caicedo<sup>†</sup>, John R. Stinchcombe<sup>‡</sup>, Kenneth M. Olsen<sup>†</sup>, Johanna Schmitt<sup>‡</sup>, and Michael D. Purugganan<sup>†§</sup>

<sup>†</sup>Genetics Department, North Carolina State University, Raleigh, NC 27695; and <sup>‡</sup>Department of Ecology and Evolutionary Biology, Brown University, Providence, RI 02912

Edited by Susan R. Wessler, University of Georgia, Athens, GA, and approved September 23, 2004 (received for review August 24, 2004)

Epistatic gene interactions are believed to be a major factor in the genetic architecture of evolutionary diversification. In *Arabidopsis thaliana*, the *FRI* and *FLC* genes mechanistically interact to control flowering time, and here we show that this epistatic interaction also contributes to a latitudinal cline in this life history trait within the species. Two major *FLC* haplogroups (*FLC<sup>A</sup>* and *FLC<sup>B</sup>*) are associated with flowering time variation in *A. thaliana* in field conditions, but only in the presence of putatively functional *FRI* alleles. Significant differences in latitudinal distribution of *FLC* haplogroups in Eurasia and North Africa also depend on the *FRI* genotype. There is significant linkage disequilibrium between *FRI* and *FLC* despite their location on separate chromosomes. Although no nonsynonymous polymorphisms differentiate *FLC<sup>A</sup>* and *FLC<sup>B</sup>*, vernalization induces the expression of an alternatively spliced *FLC* transcript that encodes a variant protein with a radical amino acid replacement associated with the two *FLC* haplogroups. Numerous polymorphisms differentiating the *FLC* haplogroups also occur in intronic regions implicated in the regulation of *FLC* expression. The features of the regulatory gene interaction between *FRI* and *FLC* in contributing to the latitudinal cline in *A. thaliana* flowering time are consistent with the maintenance of this interaction by epistatic selection. These results suggest that developmental genetic pathways and networks provide the molecular basis for epistasis, contributing to ecologically important phenotypic variation in natural populations and to the process of evolutionary diversification.

rosette leaf number | alternative splicing | linkage disequilibrium | epistatic selection | candidate gene association

Epistasis has been an enigmatic factor in the conceptualization of the evolutionary process (1, 2). Fisher (1, 3) contended that epistatic gene interactions were of limited relevance to evolutionary diversification, whereas Wright (4) argued for the central role of epistasis in evolution. Recently, renewed interest in the Wrightian view of evolutionary diversification has led to attempts to define and characterize epistasis in a wide range of micro- and macro-evolutionary phenomena. Gene interactions are now widely believed to be crucial features of the maintenance of genetic variation (5), the origins of sex and recombination (6), and the process of speciation (7). The importance of epistasis has also been bolstered by advances in quantitative trait loci (QTL) mapping that reveals examples of epistatic interactions underlying natural phenotypic variation in organisms such as *Drosophila* (8), *Arabidopsis* (5, 9), and *Caenorhabditis elegans* (10).

Concurrent with interest in the evolutionary role of epistasis, molecular geneticists have begun to identify genetic pathways and networks involved in the expression of many phenotypic features (11). These pathways provide frameworks for dissecting the molecular basis of gene interactions underlying ecologically and evolutionarily relevant phenotypic variation. Regulatory genetic pathways that control developmental processes are particularly intriguing, because it is thought that evolution of these regulatory loci underlies key aspects of evolutionary diversification (12).

Despite progress in the characterization of molecular genetic pathways, the study of epistasis continues to rely heavily on characterizing statistical interaction terms in quantitative genetic analysis (13) or defining broad chromosomal regions that contribute nonadditively to phenotypic variation (5). There have been few examples of epistasis resolved at the level of molecular interactions of naturally occurring alleles of specific genes (14–16) and no examples of regulatory gene interactions leading to natural phenotypic variation in ecologically relevant traits. Identifying regulatory gene interactions and determining their molecular mechanisms should provide key insights into the origins and importance of epistasis in evolutionary change.

Here we report on a specific regulatory gene interaction that contributes to natural variation in the timing of flowering in *Arabidopsis thaliana*. In *A. thaliana*, flowering time has been associated with size at reproduction and fecundity (17–19). Flowering time is sensitive to climatic signals, including day length (photoperiod) and prolonged cold treatment (vernalization), which serve as ecological cues to ensure that reproductive effort occurs in optimal seasonal environments (20). These signals vary systematically with latitude, and adaptation to these ecological cues would be expected to lead to latitudinal clines in the timing of flowering. Due to its distribution over a wide geographic range, latitude-dependent variation in flowering time in *A. thaliana* might be expected to be associated with molecular polymorphisms in genes that regulate the transition to flowering in response to environmental cues (21–23).

A large number of regulatory genes belonging to several interacting pathways that control flowering time in *A. thaliana* have been identified (20, 24). Genes in the vernalization pathway, including *FRIGIDA* (*FRI*) (25) and *FLC* (26, 27) (Fig. 1), condition *A. thaliana*'s response to cold temperatures associated with winter, an environmental cue that promotes flowering in the spring. A recent study has demonstrated the presence of a latitudinal cline in flowering time modulated by the *FRI* gene in *A. thaliana* (28). *FRI* encodes a 609-aa coiled-coil protein that represses the transition to flowering, and mutants of this gene display early flowering under long-day controlled-temperature conditions, as well as insensitivity to vernalization (25). Several naturally occurring null *FRI* alleles have been described, and molecular analyses indicate that these independently derived alleles result from several deletions in the coding region (25, 29).

In a common garden experiment carried out under over-winter field conditions, a latitudinal cline in bolting time is evident only in ecotypes that are homozygous for a putative functional *FRI* allele; southern ecotypes in this genotypic class flower significantly earlier than those at northern latitudes (28). In contrast, ecotypes homozy-


This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: RLN, rosette leaf number at bolting; SNP, single-nucleotide polymorphism.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. AY769347–AY769360, AY769865, and AY772670–AY772672).

<sup>§</sup>To whom correspondence should be addressed. E-mail: michael.purugganan@ncsu.edu.

© 2004 by The National Academy of Sciences of the USA



**Fig. 1.** Schematic representation of genetic pathways of flowering time in *A. thaliana*. Lines with arrows denote up-regulation of gene expression; lines with bars denote repression of gene expression.

gous for a deletion allele of *FRI* (*FRI*<sup>Δ</sup>) flower later under natural over-winter conditions and do not exhibit the cline in flowering time. Thus, it is not latitudinal divergence in *FRI* functionality that is responsible for establishing the latitudinal cline in *A. thaliana* flowering time (28). This suggests that functional divergence in another gene, epistatic to *FRI*, may be responsible for establishing this life history cline.

A possible candidate for this interacting gene is *FLC* (Fig. 1), which encodes a MADS-box transcriptional activator that represses flowering. Mutants at *FLC* suppress the late-flowering phenotype of *FRI* under nonvernalization conditions (30), and molecular analyses indicate that transcription of *FLC* is up-regulated by *FRI* (26, 27). Moreover, *FLC* expression is down-regulated by vernalization, providing a mechanism to ensure flowering after the prolonged cold of winter (26, 27). Our evolutionary genetic analysis of *FLC* reveals that nucleotide polymorphism in this gene is indeed associated with flowering time variation in *A. thaliana* and contributes to the latitudinal cline in this trait. The association of *FLC* variation with flowering time is observed only in epistatic interaction with the upstream regulatory gene *FRI*. These results demonstrate the importance of epistasis among regulatory genes in generating phenotypic variation in a key life history trait in *A. thaliana*.

## Materials and Methods

**Molecular Population Genetic Analyses.** *A. thaliana* ecotypes were obtained from the Arabidopsis Biological Resource Center (Table 2, which is published as supporting information on the PNAS web site). *Arabidopsis lyrata* seed was provided by O. Savolainen (University of Oulu, Oulu, Finland). Genomic DNA was isolated from young leaves by using a Plant DNeasy Mini kit (Qiagen, Chatsworth, CA). The complete *FLC* gene, as well as an ≈500-bp upstream region (392 bp of the promoter and the complete 5' UTR) and ≈34 bp of the 3' UTR, was amplified in overlapping fragments for 13 ecotypes of *A. thaliana* with *Taq* polymerase (Table 3, which is published as supporting information on the PNAS web site). *A. lyrata* PCRs were carried out on two partially overlapping portions of the gene by using ExTaq DNA polymerase (Takara Shuzo, Kyoto).

*A. thaliana* PCR products were purified by using QIAquick gel extraction kits (Qiagen) and sequenced directly by using cycle sequencing with BigDye terminators (Applied Biosystems). *A. lyrata* PCR products were cloned with the TOPO TA PCR cloning kit (Invitrogen); plasmid DNA from three to five independent clones per amplified region was sequenced. Sequencing was carried out at the North Carolina State University Genome Research Laboratory with a Prism 3700 automated sequencer (Applied Biosystems).

All sequences were aligned against the published Col-0 *FLC* sequence. Molecular population genetic analyses were conducted by using DNASP 3.99 (31). Haplotype trees were constructed by using a maximum parsimony analysis (branch and bound search, stepwise addition) in PAUP\* (32). Insertion/deletion polymorphisms (indels)

that were part of long mono- or dinucleotide repeats were excluded from all analyses. Other indels were included in the parsimony analysis of *A. thaliana* sequences, with each indel treated as a single character. Parsimony analyses with *A. lyrata* excluded all indels.

**Haplotype Tagging and Association Tests.** Single-nucleotide polymorphisms (SNPs) distinguishing two major *FLC* haplotype groups were identified from *A. thaliana* sequences. Genotyping of two SNPs was carried out by using dCAPS (33) in 353 ecotypes (Table 4, which is published as supporting information on the PNAS web site). Previously identified transposon insertions as well as a Col-0 repeat (34, 35) were also genotyped. Association analyses of flowering time under unvernallized growth chamber (long day and short day) and vernalized field conditions was conducted by using SAS (Version 9.1, SAS Institute, Cary, NC). Cryptic population structure (stratification) within the sample was controlled by using 92 ecotypes belonging to an unstratified population (36) (see *Supporting Text* and Fig. 6, which are published as supporting information on the PNAS web site). Sample size in association tests varied slightly depending on ecotypes for which genotypic and phenotypic data were obtained. Conditions and phenotypes measured under growth chamber and over-wintered field conditions are described in refs. 28 and 36.


**FLC Expression Analyses.** Four *A. thaliana* ecotypes with nondeletion *FRI* alleles (Est-0, Pa-1, Le-0, and Po-0) were used to evaluate *FLC* expression. Seeds were stratified at 4°C for 3 days and planted in a randomized design across two blocks. Planted seeds were placed at 20°C, 12 h per day for 7 days until germination. A block was then moved to 4°C, 12 h per day, whereas the other block remained at 20°C. Two replicates of whole-plant tissue per ecotype were collected from newly germinated plants (V0) and plants grown continuously at 4°C (V15) and 21°C (UV15) for 15 days. Each V0 replicate contained tissue from 10 plants; all others contained tissue from four plants.

RNA was extracted by using an RNeasy kit (Qiagen) and treated with DNA-free DNase (Ambion, Austin, TX). Complementary DNA was synthesized from ≈1 μg of total RNA with the Retroscript reverse transcription kit (Ambion). RT-PCR was performed by using previously described *FLC* primers (37). Product band intensity in ethidium-stained agarose gels was used as a semiquantitative measure of expression level. Details of the RT-PCR procedure and analysis are included in *Supporting Text*.

## Results

**Nucleotide Polymorphisms at the *FLC* Gene.** The latitudinal cline in *A. thaliana* flowering time modulated by putative functional *FRI* alleles may arise as a result of the activity of its downstream regulatory target gene, *FLC*. We thus examined *FLC* in 13 *A. thaliana* ecotypes and Col-0 for the presence of genetic variation. Approximately 6.2–7.4 kb were sequenced (length variation is due to the presence of indels) and include the entire coding unit, the 5' UTR, and portions of the 3' UTR and the promoter. Sixty-five SNPs and 11 indels were observed across the entire gene (Fig. 2). The silent site nucleotide diversity,  $\pi$ , is 0.004 (Table 1), which is slightly less than the mean of 0.007 observed for previously studied *A. thaliana* nuclear genes (38).

There are 11 distinct *FLC* haplotypes in our sample, and genealogical relationships among them are shown in Fig. 3. Like several *A. thaliana* genes (38), *FLC* exhibits dimorphism; haplotypes are organized into two groups (haplogroups) differentiated by 22 SNPs and six indels (Fig. 3). We refer to these haplogroups as *FLC*<sup>A</sup> and *FLC*<sup>B</sup>. All but one of these differentiating polymorphisms are in the large ≈3.5-kb first intron of the gene. Using *A. lyrata* *FLC* as an outgroup roots the tree between the *FLC*<sup>A</sup> and *FLC*<sup>B</sup> clades (Fig. 7, which is published as supporting information on the PNAS web site). No exon polymorphisms distinguish *FLC*<sup>A</sup> and *FLC*<sup>B</sup>, and the protein sequences encoded by the two haplogroups are identical.



**Fig. 2.** Polymorphisms observed in the sequenced *FLC* region. Site numbering is based on our alignment and includes indels. Polymorphisms in mononucleotide repeats longer than 4-bp and long dinucleotide repeats are not included. Positions highlighted in gray, along with the Da (1)-12 type insert (not shown), were used in genotyping.

The only nonsynonymous polymorphism in the sample occurs within *FLC<sup>A</sup>* (a conservative R to K replacement found in *Ler-0*). The two *FLC* haplogroups differ substantially in the amounts of polymorphism. The eight *FLC<sup>A</sup>* alleles have 41 nucleotide polymorphisms, with  $\pi = 0.0022$ . In contrast, the six *FLC<sup>B</sup>* alleles have seven SNPs and a 5-fold lower level of nucleotide diversity ( $\pi = 0.0004$ ) (Table 1). Patterns with an unbalanced distribution of polymorphism have sometimes been ascribed to partial selective sweeps in other biological systems (39).

We genotyped a set of 353 primarily Eurasian ecotypes for *FLC<sup>A</sup>* and *FLC<sup>B</sup>* alleles (Table 2). The *FLC<sup>A</sup>* allele is present at slightly higher frequency (57%) than *FLC<sup>B</sup>* (43%). Tajima's *D* for *FLC* is positive ( $D = 0.8579$ ), consistent with the presence of two haplotype groups at intermediate frequencies, although it does not differ significantly from predictions of the neutral-equilibrium theory. Previous studies have identified two large transposable element insertions in the first intron of *FLC* that reduce expression and lead to early flowering (34, 35). These insertions occur at low frequency in our ecotype sample (<3%) and are all nested within the *FLC<sup>A</sup>* haplogroup (Fig. 3 and Table 2). A small direct repeat insertion in intron 1, characteristic of the Col-0 ecotype (34, 35), is also nested within *FLC<sup>A</sup>* and occurs at moderately low frequency (14.7%).

**Epistasis Between *FRI* and *FLC* Is Associated with Flowering Time Variation.** Field experiments have revealed latitudinal variation in *A. thaliana* flowering time only in accessions that do not contain inactivating deletions at the *FRI* gene (28). Molecular genetic studies indicate that *FRI* up-regulates *FLC* expression (26, 27); we would thus expect that variation in the latter would show significant association with flowering time, but only in ecotypes that have an active *FRI* allele.

We tested for effects of *FRI* and *FLC* on flowering time using a set of 92 ecotypes shown to behave as an unstratified population appropriate for testing phenotypic associations (36). We genotyped these accessions for *FLC<sup>A</sup>* and *FLC<sup>B</sup>* alleles and for the two intron 1 transposable element insertions (34, 35); eight ecotypes with

transposon insertions were excluded from the analyses, as were five others for which the presence/absence of insertions could not be determined. Together with *FRI* genotypes classes previously observed for 72 of the remaining ecotypes (28), we determined the two-locus *FLC* and *FRI* deletion (*FRI<sup>Δ</sup>*) and nondeletion (*FRI*) genotypes and tested for associations with rosette leaf number at bolting (RLN) and days until bolting, both common indicators of flowering time. There is substantial broad sense heritability ( $H^2$ ) for flowering time under growth chamber and field conditions (36). RLN and days until bolting are highly genetically correlated traits (0.94) (9), and results obtained for both measures were largely congruent. We present the more conservative RLN-based results in the text; results obtained with bolting days are presented in Table 5, which is published as supporting information on the PNAS web site. Only results significant for both traits are discussed.

We did not find a significant association of *FRI* or *FLC* genotypes with flowering time under short-day controlled environmental conditions (Fig. 4A and Table 6, which is published as supporting information on the PNAS web site). We did find an association of *FRI* genotype with RLN under long-day controlled environmental conditions (two-way ANOVA:  $P = 0.0014$ ); ecotypes with *FRI* deletion alleles exhibit early flowering, as expected (Fig. 4B and Table 6). No association was seen for RLN with *FLC* or *FRI* *FLC* genotype under long-day conditions. We did, however, find significant associations for both genes and RLN under field conditions (*FRI* genotype  $P = 0.028$ ; *FLC* haplogroup  $P = 0.0049$ ), as well as for the *FRI*  $\times$  *FLC* interaction ( $P = 0.045$ ) (Fig. 4C and D and Table 6). This suggests that both genes have an effect on flowering time under ecological field conditions; examination of the variance components reveals that substantial amounts of variance in RLN are explained by *FLC*, *FRI*, and their epistatic interaction (11.1%, 4.35%, and 13.43%, respectively; Table 7, which is published as supporting information on the PNAS web site).

Post hoc tests reveal that, under field conditions, there is no significant difference in mean RLN for ecotypes with a *FRI* deletion allele (*FRI<sup>Δ</sup>* *FLC<sup>A</sup>* mean RLN =  $27.51 \pm 0.85$ ,  $n = 18$ ;

**Table 1. Features of *FLC* sequence variation**

Clade	<i>n</i>	Sequence length, bp		<i>S</i> <sup>‡</sup>	<i>A</i> <sup>§</sup>	$\pi$ <sup>  </sup>	$\theta_w$ <sup>  </sup>	Tajima's <i>D</i>
		Alignment*†	Adjusted†					
<i>FLC</i>	14	7,460	5,994	65	11	0.0044	0.0036	0.8579
<i>FLC<sup>A</sup></i>	8	7,460	6,017	41	8	0.0022	0.0028	-1.1699
<i>FLC<sup>B</sup></i>	6	7,460	6,107	7	3	0.0004	0.0005	-1.3903


\*Alignment length.

†Length excluding indels and missing data.

‡Number of segregating sites.

§Haplotype number.

||Estimates based on silent sites.



**Fig. 3.** *FLC* haplotype network. The tree represents one of four most-parsimonious arrangements; relationships among major haplotype groups do not change with alternative trees. Each line represents a single mutational step corresponding to a SNP or an indel. Mutations marked with lower-case letters are homoplasious. Characters 1919 and 7207, which are highly homoplasious (consistency index  $\leq 0.333$ ), were excluded from the tree. Shaded boxes indicate haplogroups. Haplogroup and haplotype frequencies are based on genotyping 353 *A. thaliana* accessions.


*FRI* $\Delta$  *FLC*<sup>B</sup> mean RLN =  $28.65 \pm 0.76$ ,  $n = 28$ ) (Fig. 4C). In contrast, mean RLN for accessions with a *FRI* *FLC*<sup>A</sup> genotype ( $21.95 \pm 2.35$ ,  $n = 8$ ) is significantly lower than those that are *FRI* *FLC*<sup>B</sup> ( $28.84 \pm 1.86$ ,  $n = 9$ ;  $P = 0.0049$ ). *FRI* *FLC*<sup>A</sup> ecotypes also flower significantly earlier than *FRI* $\Delta$  *FLC*<sup>A</sup> or *FRI* $\Delta$  *FLC*<sup>B</sup> ecotypes ( $P = 0.0054$  and  $P = 0.0005$ , respectively). This indicates that most genotypes flower at approximately the same time in an over-winter field environment, except for the earlier flowering of accessions with an *FLC*<sup>A</sup> allele in a *FRI* nondeletion background. Thus, an

association of *FLC* allele class with flowering time is detectable only in a functional *FRI* background.


**Linkage Disequilibrium Between *FLC* and *FRI*.** The *FRI* and *FLC* genes are not physically linked; *FRI* is located at the top of chromosome IV, whereas *FLC* is at the top of chromosome V. Despite their location on different chromosomes, flowering time variation associated with the functional interaction of *FRI* and *FLC* could lead to linkage disequilibrium between these two loci. To test for linkage disequilibrium, we obtained *FRI* genotypes for 321 ecotypes (C. A. Mays and M.D.P., unpublished data) and combined these with observed *FLC* genotypes. The frequencies of ecotypes with *FRI* $\Delta$  *FLC*<sup>A</sup> (29%) and *FRI* $\Delta$  *FLC*<sup>B</sup> (35%) are almost equivalent, indicating that *FLC* alleles are equally distributed in a *FRI* deletion allele background. The frequency of *FRI* *FLC*<sup>A</sup> (27%) is also similar to that of the former two genotypic classes. In contrast, there is a lower number of accessions with a *FRI* *FLC*<sup>B</sup> genotype (9%). Comparison with expected genotypic frequencies reveals that genotypes *FRI* $\Delta$  *FLC*<sup>B</sup> and *FRI* *FLC*<sup>A</sup> are overrepresented in our sample, whereas genotypes *FRI* $\Delta$  *FLC*<sup>A</sup> and *FRI* *FLC*<sup>B</sup> are underrepresented. This nonrandom distribution of allelic genotypes is significant in an exact test of gametic disequilibrium ( $P < 4.7 \times 10^{-7}$ ).

***FLC* Genotypes Display Latitudinal Differentiation.** The geographic distribution of *FRI* *FLC* genotypes was examined for 298 Eurasian *A. thaliana* accessions, for which we had site-of-origin information; we tested for *FRI* and *FLC* associations with latitude via a two-way analysis of covariance (ANCOVA), including longitude as a covariate. We found it necessary to control for longitude, because it is correlated with latitude in our dataset. There is a significant association of *FLC* and *FRI* *FLC* genotype with latitude ( $P = 0.0025$  and  $P = 0.001$ , respectively; Table 8, which is published as supporting information on the PNAS web site). Post hoc tests reveal significant difference in latitudinal distribution of *FLC*<sup>A</sup> and *FLC*<sup>B</sup> alleles in a *FRI* nondeletion background ( $P < 0.001$ ) but not in a *FRI* $\Delta$  background. Plants that have the *FRI* *FLC*<sup>B</sup> genotype are found primarily in northern latitudes (mean latitude =  $51.76^\circ\text{N} \pm 0.55$ ,  $n = 24$ ), whereas accessions with *FRI* *FLC*<sup>A</sup> genotypes have a more southern distribution (mean latitude =  $48.37^\circ\text{N} \pm 0.56$ ,  $n = 78$ ). In contrast, the mean latitudes for *FRI* $\Delta$  *FLC*<sup>A</sup> ( $50.16^\circ\text{N} \pm 0.36$ ,  $n = 89$ ) and *FRI* $\Delta$  *FLC*<sup>B</sup> ( $50.26^\circ\text{N} \pm 0.26$ ,  $n = 107$ ) are almost equal. To determine whether the geographic differentiation of *FRI* *FLC*<sup>A</sup> and *FRI* *FLC*<sup>B</sup> genotypes is a consequence of population structure within *A. thaliana*, we examined the distribution of genotypes in our unstratified dataset. As before, the associations of *FLC* haplogroup and *FRI* *FLC* genotype with latitude are significant ( $P = 0.013$  and  $P = 0.038$ , respectively; Table 8). There is significant latitudinal differentiation between *FRI* *FLC*<sup>A</sup> and *FRI* *FLC*<sup>B</sup> genotypes ( $P = 0.022$ ) and no differentiation between *FRI* $\Delta$  *FLC*<sup>A</sup> and *FRI* $\Delta$  *FLC*<sup>B</sup> genotypes. This suggests that the distribution of *FLC* genotypes in a *FRI* nondeletion background is not random and is not a result of cryptic population structure.

The preferential northern distribution of *FRI* *FLC*<sup>B</sup> genotypes, which are later flowering under field conditions, and southern distribution of earlier-flowering *FRI* *FLC*<sup>A</sup> genotypes could explain the latitudinal cline in flowering time observed for putatively functional *FRI* genotypes (Fig. 8, which is published as supporting information on the PNAS web site). However, the associations of both genotype and flowering time with latitude could also indicate that other factors correlated with latitude are responsible for the observed variation in flowering time. We tested this hypothesis by running an analysis of covariance (ANCOVA) on our unstratified sample, including *FRI* and *FLC* genotypes as main effects and latitude as a covariate. Because flowering time in the common garden has also been shown to be significantly associated with mean January precipitation at site of origin (28), this variable was also



**Fig. 4.** Associations between *FRI* *FLC* genotype and flowering time. Bars that differ in roman numerals are significantly different at  $P < 0.006$  by Fisher's protected least-significant difference test. Error bars indicate standard error. (A) Short-day growth chamber conditions. (B) Long-day growth chamber conditions. (C and D) Vernalized over-wintering field conditions.



**Fig. 5.** Alternative splicing at *FLC*. (A) RT-PCR of *FLC* under different treatments. Plants at V0 were collected soon after germination and were not subjected to vernalization. Plants at V15 were grown at 4°C for 15 days after germination. (B) Structure of the alternative-splice *FLC* transcript. The upper gene model encompasses the region sequenced for *FLC*. White boxes represent UTRs, black boxes represent exon regions, and lines represent introns and promoter regions. Slender bars indicate splice sites of the alternative *FLC* product. The lower gene model represents the alternative transcript after splicing has occurred. (C) Predicted protein of the *FLC* alternative splice transcript, indicating the amino acid replacement polymorphism between the *FLC<sup>A</sup>* and *FLC<sup>B</sup>* haplogroups.

included as a covariate. After controlling for the two geographic variables, the effects of *FLC* and the *FRI* × *FLC* interaction on RLN remain significant ( $P = 0.0053$ ,  $P = 0.025$  respectively; Table 9, which is published as supporting information on the PNAS web site).

**Alternative Splicing at *FLC<sup>A</sup>* and *FLC<sup>B</sup>*.** There is no difference in the protein sequences encoded by *FLC<sup>A</sup>* and *FLC<sup>B</sup>* haplogroup alleles; instead, nearly all polymorphisms that differentiate these two haplogroups are found in the ≈3.5-kb intron 1 of *FLC*. Large introns in MADS-box regulatory genes are known to contain key cis-regulatory elements (40), and regions of the *FLC* first intron have been implicated in the control of *FLC* expression (41, 42). These observations suggest that associations of *FLC* haplogroups with flowering time may arise from regulatory differences between these two naturally occurring alleles.

To test this hypothesis, we conducted RT-PCR analysis of *FLC* transcripts from four ecotypes. Consistent with previous reports, a decrease in the expression levels of *FLC* is evident after 15 days of vernalization in ecotypes from both *FRI FLC<sup>A</sup>* and *FRI FLC<sup>B</sup>* genotype classes (Fig. 5A and Fig. 9, which is published as supporting information on the PNAS web site). Moreover, an additional shorter transcript is induced in a vernalization treatment of 15 days (Fig. 5A). This shorter transcript is not observed at high levels in plants grown at normal temperature for 15 days (data not shown). The shorter transcript is an alternatively spliced *FLC* transcript that uses a nonconsensus cryptic splice donor site in intron 1 (GA) and splice acceptor site in intron 6 (CT) and contains all of the first and last exon of the *FLC* gene, as well as 73 bp of intron 1 and 15 bp of intron 6 (Fig. 5B). It encodes an in-frame 124-aa protein, which includes the sequence-specific DNA-binding/dimerization MADS domain and the C-terminal region of the normal *FLC* protein, as well as a novel intervening protein sequence. The alternatively spliced transcript has an A to T transversion at codon 79 of the ORF, resulting in a Cys to Ser radical amino acid replacement between the putative *FLC<sup>A</sup>* and *FLC<sup>B</sup>* alternate protein products (Fig. 5C).

We did not detect significant differences in the levels of expression of the full-length or alternative *FLC* transcripts between the two haplogroups under the different treatments (Fig. 9). However, we did observe a tendency for a higher expression of full-length *FLC* in *FRI FLC<sup>A</sup>* ecotypes under vernalization conditions (Fig. 9B). There is also a tendency for lower levels of the alternative-splice transcript in *FRI FLC<sup>A</sup>* ecotypes, but this is partially obscured by the high levels of alternative transcript contributed by one replicate of the Pa-1 ecotype (Fig. 9C). Because the modest size of our assay may have limited power, we cannot rule out the possibility

of subtle differences in the levels of the *FLC* transcripts between *FLC* haplogroups.

## Discussion

Variation in flowering time is a key feature of the life histories of flowering plants and determines crucial aspects of the ecology of plant reproduction; thus, there has been intense interest in determining the genetic architecture of this trait. A recent study has demonstrated that a latitudinal cline in *A. thaliana* flowering time under field over-winter conditions is modulated by putative functional alleles of the regulatory gene *FRI* (28). Ecotypes with putatively functional *FRI* alleles located in southern Eurasian latitudes flower significantly earlier than northern ecotypes. In contrast, there is no correlation of flowering time with latitude among ecotypes that have deletion alleles of *FRI*. These results suggest a genetic model in which the latitudinal cline in *A. thaliana* flowering time depends on another gene whose function requires an active *FRI* allele.

*FRI* and the MADS-box transcription factor *FLC* are two genes known to mechanistically interact to control the transition to flowering (26, 27). Two lines of evidence indicate that the latitudinal cline in *A. thaliana* flowering time is established, in part, by epistasis between these two regulatory genes. First, *FRI* is known to up-regulate *FLC* expression, so that ecotypes with deletion alleles of *FRI* have reduced *FLC* activity and cannot express phenotypes associated with *FLC* variation. Our results reveal that variation in flowering time associated with *FLC* is observed only in plants that have a putative functional *FRI* allele, in accordance with this model (Fig. 4). Second, we would predict that *FLC* haplogroups display differences in latitudinal distribution only in ecotypes with putatively functional *FRI* alleles; this prediction is also borne out by our data. Together, these results demonstrate that *FLC*, in epistatic interaction with *FRI*, forms part of the molecular basis for the latitudinal cline in *A. thaliana* flowering time.

The precise molecular mechanism for differential *FLC* function associated with flowering time variation requires further investigation. Based on our results, two possible mechanisms could account for functional differences among haplogroups. One mechanism relies on our discovery that vernalization induces expression of an alternatively spliced *FLC* transcript. The alternate transcript encodes a modified MADS-domain protein, which may act as a dominant negative repressor by competing for target promoter-binding sites or by dimerizing with *FLC*. Unlike normal *FLC*, the modified proteins encoded by the *FLC<sup>A</sup>* and *FLC<sup>B</sup>* alternative transcripts differ by a radical amino acid change, potentially leading to differential activity. A second mechanism is suggested by previous studies of regions in the first intron of *FLC* involved in the regulation of *FLC* expression (41, 42). Some of these regions

contain polymorphisms differentiating *FLC<sup>A</sup>* and *FLC<sup>B</sup>*, potentially leading to differential expression. Expression analyses of *FLC<sup>A</sup>* and *FLC<sup>B</sup>* ecotypes do not reveal significant differences in the expression of either of the *FLC* transcripts; however, the small scale of the experiments might not be sufficient to reveal subtle differences in levels of expression. Both of the mechanisms for *FLC* differential function (alternate protein activity or expression differences) provide clear testable hypotheses; these studies are currently in progress.

The finding that the latitudinal cline in *A. thaliana* flowering time is associated with an epistatic regulatory gene interaction is particularly noteworthy. Although selection on epistatic interactions has been hypothesized, particularly in the maintenance of coadapted gene complexes, it is presumed that recombination would break apart allelic correlations unless selection were sufficiently strong (43). Epistatic selection has been suggested to be a significant evolutionary force among linked loci or in nonrandom mating or highly structured populations (44), although these restrictive features have limited the number of documented cases. Several lines of evidence, however, indicates that the epistatic interaction between the *FRI* and *FLC* regulatory genes is maintained by selection. First, the interaction between *FRI* and *FLC* clearly contributes to natural phenotypic variation in an ecologically important life history trait. Second, the flowering time variation caused by alternative *FLC* alleles in a *FRI* background is distributed along a latitudinal cline; latitudinal clines are classically regarded as strong evidence of selection associated with geographically structured climatic variables (45). Third, linkage disequilibrium among interacting loci has been regarded as a key signature of epistatic selection; the skewed genotypic associations between *FRI* and *FLC* alleles suggest that the allele combinations are targeted by selection. Although each of these results alone may be insufficient to conclude the action of selection, together they present a compelling case to suggest that this epistatic regulatory gene interaction is maintained by selection for flowering time variation across the species range of *A. thaliana*.

The precise nature and direction of selection on the *FRI FLC* genotypes are unknown, but several possibilities exist. The earlier

flowering of *FRI FLC<sup>A</sup>* ecotypes under over-wintering conditions suggests that plants carrying this genotype may be more sensitive to vernalization. The differential geographical distribution of *FRI FLC<sup>A</sup>* and *FRI FLC<sup>B</sup>* genotypes could be due to spatially heterogeneous selection. Sensitivity to vernalization might be favored in southern latitudes, where the vernalization cues of winter are milder; early flowering in response to vernalization could also serve as a mechanism to guarantee flowering before the summer heat (28). Alternatively, less sensitivity to vernalization might be favored in northern latitudes, where colder and longer winters could lead to premature flowering in highly vernalization-sensitive plants; later flowering may also lead to increased fitness associated with larger plant size and greater fruit production (18). The skewed frequencies of *FRI FLC* genotypes may be due to the availability of habitats where alternative vernalization-sensitive phenotypes are favored. Selection on unknown traits influenced by *FLC* haplogroups in both *FRI* backgrounds (deletion and nondeletion) may also be a factor shaping both the geographical distribution and frequencies of these genotypes.

Although the role of epistasis in evolutionary diversification has been the subject of intense debate, there are few known examples of natural phenotypic variation mediated by epistatic interactions between defined genes. Molecular genetic characterization of regulatory gene interactions, such as those observed in the *A. thaliana* flowering time network, provides evolutionary geneticists with concrete mechanistic frameworks to define epistatic interactions that underlie ecologically and evolutionarily relevant trait variation. We can now begin to bridge the gap between the population epistasis of the evolutionary geneticist and the mechanistic epistasis of molecular and developmental geneticists (46, 47) and to dissect, at the molecular level, the mechanisms and roles of gene interactions in the process of evolutionary change.

We thank S. Rusza for technical assistance and T. Korves, R. C. Moore, and C. Mays for discussion and suggestions on the manuscript. This work was funded in part by a National Science Foundation Integrated Research Challenges in Environmental Biology grant (to M.D.P., J.S., and T. F. C. Mackay).

1. Fenster, C. B., Galloway, L. F. & Chao, L. (1997) *Trends Ecol. Evol.* **12**, 282–286.
2. Wade, M. J., Winther, R. G., Agrawal, A. F. & Goodnight, C. J. (2001) *Trends Ecol. Evol.* **16**, 498–504.
3. Fisher, R. A. (1930) *The Genetical Theory of Natural Selection* (Oxford Univ. Press, Oxford).
4. Wright, S. (1931) *Genetics* **16**, 97–159.
5. Weinig, C., Dorn, L. A., Kane, N. C., German, Z. M., Halldorsdottir, S. S., Ungerer, M. C., Toyonaga, Y., Mackay, T. F., Purugganan, M. D. & Schmitt, J. (2003) *Genetics* **165**, 321–329.
6. Charlesworth, B. (1990) *Genet. Res.* **55**, 199–221.
7. Orr, H. A. (1995) *Genetics* **139**, 1805–1813.
8. Dilda, C. L. & Mackay, T. F. C. (2002) *Genetics* **162**, 1655–1674.
9. Ungerer, M. C., Halldorsdottir, S. S., Modliszewski, J. L., Mackay, T. F. C. & Purugganan, M. D. (2002) *Genetics* **160**, 1133–1151.
10. Shook, D. R. & Johnson, T. E. (1999) *Genetics* **153**, 1233–1243.
11. Cork, J. M. & Purugganan, M. D. (2004) *BioEssays* **26**, 479–484.
12. Doebley, J. & Lukens, L. (1998) *Plant Cell* **10**, 1075–1082.
13. Hard, J. J., Bradshaw, W. E. & Holzapfel, C. M. (1992) *Genetics* **131**, 389–396.
14. Rawson, P. D. & Burton, R. S. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 12955–12958.
15. Templeton, A. R. (2000) in *Epistasis and the Evolutionary Process*, eds. Wolf, J. B., Brodie, E. D., III, & Wade, M. J. (Oxford Univ. Press, New York), pp. 41–57.
16. McKechnie, S. W. & Geer, B. W. (1988) *Genet. Res.* **52**, 179–184.
17. Mitchell-Olds, T. (1996) *Evolution (Lawrence, Kans.)* **50**, 140–145.
18. Clauss, M. J. & Arssen, L. W. (1994) *J. Ecol.* **82**, 447–455.
19. Dorn, L. A., Pyle, E. H. & Schmitt, J. (2000) *Evolution (Lawrence, Kans.)* **54**, 1982–1994.
20. Mouradov, A., Cremer, F. & Coupland, G. (2002) *Plant Cell* **14**, S111–S130.
21. Nordborg, M. & Bergelson, J. (1999) *Am. J. Bot.* **86**, 470–475.
22. Stenoien, H. K., Fenster, C. B., Kuitinen, H. & Savolainen, O. (2002) *Am. J. Bot.* **89**, 1604–1608.
23. Pigliucci, M. & Marlow, E. T. (2001) *Oecologia* **127**, 501–508.
24. Simpson, G. G. & Dean, C. (2002) *Science* **296**, 285–289.
25. Johanson, U., West, J., Lister, C., Michaels, S., Amasino, R. & Dean, C. (2000) *Science* **290**, 344–347.
26. Sheldon, C. C., Burn, J. E., Perez, P. P., Metzger, J., Edwards, J. A., Peacock, W. J. & Dennis, E. S. (1999) *Plant Cell* **11**, 445–458.
27. Michaels, S. D. & Amasino, R. M. (1999) *Plant Cell* **11**, 949–956.
28. Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., Purugganan, M. D. & Schmitt, J. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 4712–4717.
29. Le Corre, V., Roux, F. & Reboud, X. (2002) *Mol. Biol. Evol.* **19**, 1261–1271.
30. Michaels, S. D. & Amasino, R. M. (2001) *Plant Cell* **13**, 935–941.
31. Rozas, J. & Rozas, R. (1999) *Bioinformatics* **15**, 174–175.
32. Swofford, D. L. (2000) PAUP\*, Phylogenetic Analysis Using Parsimony (\*and Other Methods) (Sinauer, Sunderland, MA).
33. Neff, M. M., Turk, E. & Kalishman, M. (2002) *Trends Genet.* **18**, 613–615.
34. Gazzani, S., Gendall, A. R., Lister, C. & Dean, C. (2003) *Plant Physiol.* **132**, 1107–1114.
35. Michaels, S. D., He, Y. H., Scortecci, K. C. & Amasino, R. M. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 10102–10107.
36. Olsen, K. M., Halldorsdottir, S. S., Stinchcombe, J. R., Weinig, C., Schmitt, J. & Purugganan, M. D. (2004) *Genetics* **167**, 1361–1369.
37. Michaels, S. D., Ditta, G., Gustafson-Brown, C., Pelaz, S., Yanofsky, M. & Amasino, R. M. (2003) *Plant J.* **33**, 867–874.
38. Yoshida, K., Kamiya, T., Kawabe, A. & Miyashita, N. T. (2003) *Genes Genet. Syst.* **78**, 11–21.
39. Yi, S. & Charlesworth, B. (2003) *Genet. Res.* **82**, 101–106.
40. Sieburth, L. E. & Meyerowitz, E. M. (1997) *Plant Cell* **9**, 355–365.
41. Sheldon, C. C., Conn, A. B., Dennis, E. S. & Peacock, W. J. (2002) *Plant Cell* **14**, 2527–2537.
42. He, Y. H., Michaels, S. D. & Amasino, R. M. (2003) *Science* **302**, 1751–1754.
43. Lewontin, R. (1974) *The Genetic Basis of Evolutionary Change* (Columbia Univ. Press, New York).
44. Clegg, M. T., Allard, R. W. & Kahler, A. L. (1972) *Proc. Natl. Acad. Sci. USA* **69**, 2474–2478.
45. Endler, J. A. (1986) *Natural Selection in the Wild* (Princeton Univ. Press, Princeton, NJ).
46. Brodie, E. D., III (2000) in *Epistasis and the Evolutionary Process*, eds. Wolf, J. B., Brodie, E. D., III, & Wade, M. J. (Oxford Univ. Press, New York), pp. 3–19.
47. Phillips, P. C. (1998) *Genetics* **149**, 1167–1171.